

# Human 2B4/CD244/SLAMF4 Protein

Cat. No. 2B4-HM101

## Description

<b>Source</b>	Recombinant Human 2B4/CD244/SLAMF4 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Cys22-Arg221.
<b>Accession</b>	Q9BZW8-2
<b>Molecular Weight</b>	The protein has a predicted MW of 23.1 kDa. Due to glycosylation, the protein migrates to 45-70 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

2B4 (CD244) is expressed by memory-phenotype CD8 T cells and all natural killer (NK) cells. The ligand for 2B4, CD48, is expressed on hematopoietic cells. 2B4 is conserved in humans and mice, and a number of reports have linked 2B4 with activation of lymphocytes. Engagement of 2B4 on NK cell surfaces with specific antibodies or CD48 can trigger cell mediated cytotoxicity, interferon  $\gamma$  secretion, phosphoinositol turnover and NK cell invasiveness.


## Assay Data

### Bis-Tris PAGE



Human 2B4 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC




The purity of Human 2B4 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

**Human 2B4, His Tag ELISA**

0.2µg Human 2B4, His Tag Per Well



Immobilized Human 2B4, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Human CD48, hFc Tag with the EC50 of 92.3ng/ml determined by ELISA.