

Human 2B4/CD244/SLAMF4 Protein

Cat. No. 2B4-HM101

Description

Source	Recombinant Human 2B4/CD244/SLAMF4 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Cys22-Arg221.
Accession	Q9BZW8-2
Molecular Weight	The protein has a predicted MW of 23.1 kDa. Due to glycosylation, the protein migrates to 45-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

2B4 (CD244) is expressed by memory-phenotype CD8 T cells and all natural killer (NK) cells. The ligand for 2B4, CD48, is expressed on hematopoietic cells. 2B4 is conserved in humans and mice, and a number of reports have linked 2B4 with activation of lymphocytes. Engagement of 2B4 on NK cell surfaces with specific antibodies or CD48 can trigger cell mediated cytotoxicity, interferon γ secretion, phosphoinositol turnover and NK cell invasiveness.

Assay Data


Tris-Bis PAGE



Human 2B4 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human 2B4 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human 2B4, His Tag ELISA

0.1µg Human 2B4, His Tag Per Well



Immobilized Human 2B4, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human CD48, hFc Tag with the EC50 of 0.34µg/ml determined by ELISA.