

Biotinylated Cynomolgus ACE2/ACEH Protein

Cat. No. ACE-CM401B

Description

Source	Recombinant Biotinylated Cynomolgus ACE2/ACEH Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gln18-Val739.
Accession	A0A2K5X283
Molecular Weight	The protein has a predicted MW of 86.5 kDa. Due to glycosylation, the protein migrates to 110-140 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

ACE2 (Angiotensin I Converting Enzyme 2) is a Protein Coding gene. Diseases associated with ACE2 include Severe Acute Respiratory Syndrome and Neurogenic Hypertension. The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxypeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7.

Assay Data

Bis-Tris PAGE




Biotinylated Cynomolgus ACE2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Biotinylated Cynomolgus ACE2/ACEH Protein

Cat. No. ACE-CM401B

Assay Data




The purity of Biotinylated Cynomolgus ACE2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Cynomolgus ACE2, His Tag ELISA

0.1 μ g SARS-COV-2 Spike RBD, hFc Tag Per Well



Immobilized SARS-COV-2 Spike RBD, hFc Tag at 1 μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Cynomolgus ACE2, His Tag with the EC50 of 0.28 μ g/ml determined by ELISA.