Cynomolgus ALK-7/Activin RIC Protein

Cat. No. ALK-CM107



Description	
Source	Recombinant Cynomolgus ALK-7/Activin RIC Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Gly25-Glu113.
Accession	A0A2K5VZP2
Molecular Weight	The protein has a predicted MW of 11.2 kDa. Due to glycosylation, the protein migrates to 20-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 90% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in 20mM Tris, 150mM NaCl (pH 7.5). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Activin receptor-like kinase 7 (ALK7), a member of type I transforming growth factor- β (TGF- β) receptors, is correlated with pathogenic risks of type 2 diabetes mellitus and cardiovascular diseases and may be involved in cardiovascular remodeling.


Assay Data

Bis-Tris PAGE



Cynomolgus ALK-7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Cynomolgus ALK-7 is greater than 90% as determined by SEC-HPLC.

KAGTUS


Assay Data

ELISA Data

Cynomolgus ALK-7, His Tag ELISA 0.05μg Cynomolgus ALK-7, His Tag Per Well



BLI Data



Loaded Anti-ALK-7 Antibody, hFc-Avi Tag on ProA-Biosensor can bind Cynomolgus ALK-7, His Tag with an affinity constant of 5.30 nM as determined in BLI assay (Gator® Prime).

response curve for Anti-ALK-7 Antibody, hFc Tag

Immobilized Cynomolgus ALK-7, His Tag at

 $0.5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose