

Human ALK-7/Activin RIC Protein

Cat. No. ALK-HM207

Description

Source	Recombinant Human ALK-7/Activin RIC Protein is expressed from HEK293 with hFc tag at the C-terminus. It contains Glu21-Glu113.
Accession	Q8NER5-1
Molecular Weight	The protein has a predicted MW of 35.96 kDa. Due to glycosylation, the protein migrates to 48-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1EU per µg by the LAL method.
Purity	>95% as determined by Bis-Tris PAGE >95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Arterial stiffness is an important feature of diabetic macrovascular complications. Activin receptor-like kinase 7 (ALK7), a member of type I transforming growth factor-β (TGF-β) receptors, is correlated with pathogenic risks of type 2 diabetes mellitus and cardiovascular diseases and may be involved in cardiovascular remodeling.


Assay Data

Bis-Tris PAGE



Human ALK-7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human ALK-7 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human ALK-7, hFc Tag ELISA
0.1µg Human ALK-7, hFc Tag Per Well



Immobilized Human ALK-7, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-ALK-7 Antibody, hFc Tag with the EC50 of 4.1ng/ml determined by ELISA (QC Test).