Human AMHRII Protein

Cat. No. AMH-HM0R2



Description	
Source	Recombinant Human AMHRII Protein is expressed from HEK293 without tag.
	It contains Pro18-Ser144.
Accession	Q16671-1
Molecular Weight	The protein has a predicted MW of 14.3 kDa. Due to glycosylation, the protein migrates to 30-40 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

Background


Storage

-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3-6 months after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

The aim of the current study was to explore whether anti-Müllerian hormone receptor II (AMHRII) genetic variants influence the hormonal profile and the ovarian response to standard gonadotropin stimulation of women undergoing medically assisted reproduction. Three hundred in vitro fertilization or intracytoplasmic sperm injection patients constituted the study population, while 300 women with at least one spontaneous pregnancy participated as controls.


Assay Data

Tris-Bis PAGE



Human AMHRII on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human AMHRII is greater than 95% as determined by SEC-HPLC.

Human AMHRII Protein

Cat. No. AMH-HM0R2




Assay Data

ELISA Data

Human AMHRII, No Tag ELISA

 $0.1 \mu g$ Human AMHRII, No Tag Per Well



Immobilized Human AMHRII at 1µg/ml (100µl/Well) on the plate. Dose response curve for Anti-AMHRII Antibody, hFc Tag with the EC50 of 15.6ng/ml determined by ELISA.