

Cynomolgus ANGPTL4/Angiopoietin-like 4 Protein

Cat. No. ANG-CM1L4

Description

Source	Recombinant Cynomolgus ANGPTL4/Angiopoietin-like 4 Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Arg164-Ser406.
Accession	XP_045236104.1
Molecular Weight	The protein has a predicted MW of 28.46 kDa. Due to glycosylation, the protein migrates to 35-40 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Candidates for this common regulatory system include signals mediated by peroxisome proliferator-activated regulator and its response factor, angiopoietin-like 4. The expression and bioactivity of angiopoietin-like 4, an adipocytokine that was originally reported to have an angiogenic function, have been detected not only in the vascular system and adipose tissue but also in rheumatoid joints.

Assay Data


Tris-Bis PAGE



Cynomolgus ANGPTL4 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Cynomolgus ANGPTL4 is greater than 95% as determined by SEC-HPLC.