Human CD13/ANPEP Protein, Ultra Low Endotoxin





Description	
Source	Recombinant Human CD13/ANPEP Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Lys69-Lys967.
Accession	P15144
Molecular Weight	The protein has a predicted MW of 103.8 kDa. Due to glycosylation, the protein migrates to 115-130 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.001 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD13/aminopeptidase N is a widely expressed ectoenzyme with multiple functions. As an enzyme, CD13 regulates activities of numerous cytokines by cleaving their N-terminals and is involved in Ag processing by trimming the peptides bound to MHC class II. Independent of its enzymatic activity, cell membrane CD13 functions by cross-linking-induced signal transduction, regulation of receptor recycling, enhancement of FcyR-mediated phagocytosis, and acting as a receptor for cytokines.


Assay Data

Bis-Tris PAGE



Human CD13 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human CD13 is greater than 95% as determined by SEC-HPLC.

Human CD13/ANPEP Protein, Ultra Low Endotoxin

Cat. No. ANP-HM101-UL

KAGTUS

Assay Data

Bioactivity Data

Measured by its ability to cleave the fluorogenic peptide substrate, Ala-7-amido-4-methylcoumarin (Ala-AMC). The specific activity is >2500 pmol/min/ μ g.