

FITC-Labeled Human ANGPT2/Angiopoietin-2 Protein

Cat. No. APN-HM401F

Description

Source	Recombinant FITC-Labeled Human ANGPT2/Angiopoietin-2 Protein is expressed from HEK293 with His tag and Avi tag at the N-terminus. It contains Lys275-Phe496.
Accession	O15123-1
Molecular Weight	The protein has a predicted MW of 28.4 kDa. Due to glycosylation, the protein migrates to 30-40 kDa based on Bis-Tris PAGE result.
Wavelength	Excitation Wavelength: 490 nm Emission Wavelength: 520 nm
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	>95% as determined by Tris-Bis PAGE

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Angiopoietin-2 (Ang-2; also ANGPT2) is a secreted glycoprotein that plays a complex role in angiogenesis and inflammation. Mature Ang-2 is 478 amino acids in length. Ang2 is widely expressed during development, but it is restricted postnatally to highly angiogenic tissues such as the placenta, ovaries, and uterus. It is particularly abundant in vascular endothelial cells (EC) where it is stored in intracellular Weibel Palade bodies..

Assay Data

Bis-Tris PAGE



FITC-Labeled Human ANGPT2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.