
# Biotinylated Cynomolgus APRIL/TNFSF13 Trimer Protein

#### Cat. No. APR-CM410B

Description	
Source	Recombinant Biotinylated Cynomolgus APRIL/TNFSF13 Trimer Protein is expressed from HEK293 with His tag, Flag tag and Avi tag at the N-Terminus.
	It contains Lys112-Leu250 trimer design.
Accession	A0A2K5TJA1
Molecular Weight	The protein has a predicted MW of 52.1 kDa. Due to glycosylation, the protein migrates to 56-65 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, 200 mM L-arginine (pH 8.0). Normally 8% mannitol is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The APRIL (a proliferation-inducing ligand), also known as TNFSF13, TALL2, TRDL1, and CD256, is a member of the TNF ligand superfamily.Both APRIL and its close relative BAFF bind and signal through the TNF superfamily receptors TACI and BCMA, while BAFF additionally functions through BAFFR.

### Assay Data

**Bis-Tris PAGE** 



Biotinylated Cynomolgus APRIL Trimer on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


KAGTUS

## Biotinylated Cynomolgus APRIL/TNFSF13 Trimer Protein

## Cat. No. APR-CM410B

#### Assay Data






The purity of Biotinylated Cynomolgus APRIL Trimer is greater than 95% as determined by SEC-HPLC.

#### ELISA Data







Immobilized Biotinylated Cynomolgus APRIL (Trimer), His-Avi Tag at 1µg/ml (100µl/Well) on streptavidin (5µg/ml) precoated plate. Dose response curve for Cynomolgus BCMA, hFc Tag with the EC50 of 2.0ng/ml determined by ELISA.