

Rat Activin RIIB/ACVR2B Protein, Ultra Low Endotoxin

Cat. No. ARB-RM12B-UL

Description

Source	Recombinant Rat Activin RIIB/ACVR2B Protein is expressed from HEK293 with His tag at the C-terminus. It contains Ser19-Thr137.
Accession	P38445
Molecular Weight	The protein has a predicted MW of 15.29 kDa. Due to glycosylation, the protein migrates to 30-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

ActRIIB (activin receptor type-2B) is an activin receptor subtype constitutively expressed in the whole body, playing a role in cellular proliferation, differentiation, and metabolism. For its various physiological activities, ActRIIB interacts with activin and multiple other ligands including myostatin (MSTN), growth differentiation factor 11 (GDF11), and bone morphogenetic protein 9 (BMP9).

Assay Data


Bis-Tris PAGE



Rat Activin RIIB on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Rat Activin RIIB is greater than 95% as determined by SEC-HPLC.

ELISA Data

Rat Activin RIIB, His Tag ELISA

0.1µg Human Activin A, No Tag Per Well



Immobilized Human Activin A, No Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Rat Activin RIIB, His Tag with the EC50 of 12.7ng/ml determined by ELISA (QC Test).