

Human ASGR1 Protein

Cat. No. ASG-HM1R1



Description

Source	Recombinant Human ASGR1 Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Gln62-Leu291.
Accession	P07306-1
Molecular Weight	The protein has a predicted MW of 27.49 kDa. Due to glycosylation, the protein migrates to 38-55 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

ASGR1 is a liver-specific surface marker that is exclusively expressed in liver and mediates internalization and lysosomal degradation of blood asialoglycoproteins.


Assay Data

Tris-Bis PAGE



Human ASGR1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human ASGR1 is greater than 95% as determined by SEC-HPLC.