### **Biotinylated Human AXL Protein**

#### Cat. No. AXL-HM401B



Description	
Source	Recombinant Biotinylated Human AXL Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Ala26-Pro449.
Accession	P30530-1
Molecular Weight	The protein has a predicted MW of 48.7 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

### Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### **Background**

Axl, a member of the TAM (Tyro3, Axl, Mer) family, and its inhibitors can specifically break the kinase signaling nodes, allowing advanced patients to regain drug sensitivity with improved therapeutic efficacy. Overexpression and activation of Axl receptor tyrosine kinase have been widely accepted to promote cell proliferation, chemotherapy resistance, invasion, and metastasis in several human cancers, such as lung, breast, and pancreatic cancers.


### **Assay Data**

#### **Bis-Tris PAGE**



Biotinylated Human AXL on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### **SEC-HPLC**



The purity of Biotinylated Human AXL is greater than 95% as determined by SEC-HPLC.

## Biotinylated Human AXL Protein


Cat. No. AXL-HM401B



### **Assay Data**

## **ELISA Data**

# **Biotinylated Human Axl, His Tag ELISA** 0.2μg Biotinylated Human Axl, His Tag Per Well



Immobilized Biotinylated Biotinylated Human Axl, His Tag at 2µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-AXL Antibody, hFc Tag with the EC50 of 13.0ng/ml determined by ELISA.