
Human B7-2/CD86 Protein

Cat. No. B72-HM286

ϗͶϲϿ·Ͷϩ


Description		
Decembinent Luncer DZ 2/2020 Protein is surroused from LIEK202 with hEster at the		
Source Recombinant Human B7-2/CD86 Protein is expressed from HEK293 with hFc tag at the	C-Terminus.	
It contains Leu26-Pro247.	It contains Leu26-Pro247.	
Accession P42081-1		
Molecular WeightThe protein has a predicted MW of 52 kDa. Due to glycosylation, the protein migrates to Tris PAGE result.	75-90 kDa based on Bis-	
Endotoxin Less than 1EU per µg by the LAL method.		
Purity > 95% as determined by Bis-Tris PAGE		
> 95% as determined by HPLC		
Formulation and Storage		
Formulation Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is adder lyophilization.	ed as protectant before	
Reconstitution Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ Dissolve the lyophilized protein in distilled water.	/ml is recommended.	
Storage -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after receipt to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-the storage in the protein into smaller quantities for optimal storage.		
Background		
B7-1 and B7-2 are homologous costimulatory ligands expressed on the surface of antige (APCs). Binding of these molecules to the T cell costimulatory receptors, CD28 and CTL activation and regulation of T cell immunity. B7-1 and B7-2 do not form hetero-oligomers biological relevance of dimeric and monomeric state of B7-1 and B7-2, respectively.	A-4, is essential for the	
Assay Data		

Bis-Tris PAGE



Human B7-2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human B7-2 is greater than 95% as determined by SEC-HPLC.


Human B7-2/CD86 Protein

Cat. No. B72-HM286

Assay Data

ELISA Data

Human B7-2, hFc Tag ELISA 0.5µg Human CTLA-4, His Tag Per Well



Immobilized Human CTLA-4, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Human B7-2, hFc Tag with the EC50 of 125.7ng/ml determined by ELISA.

ϗνωτυς