

Human BAFF/TNFSF13B/CD257 Trimer Protein, Ultra Low Endotoxin



Cat. No. BAF-HM213-UL


Description	
Source	Recombinant Human BAFF/TNFSF13B/CD257 Trimer Protein is expressed from HEK293 with monomeric hFc tag and Flag tag at the N-Terminus. It contains Thr141-Leu285 trimer design.
Accession	Q9Y275-1
Molecular Weight	The protein has a predicted MW of 74.9 kDa same as Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
B-cell activating factor (BAFF) also known as tumor necrosis factor ligand superfamily member 13B is a protein that in humans is encoded by the TNFSF13B gene. BAFF is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFF-R.	


Assay Data

Bis-Tris PAGE



Human BAFF (Trimer) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC



The purity of Human BAFF (Trimer) is greater than 95% as determined by SEC-HPLC.


Assay Data

ELISA Data



Immobilized Human BAFF Trimer, hFc Tag at 5µg/ml (100µl/Well) on the plate. Dose response curve for Human BAFFR, His Tag with the EC50 of 4.7ng/ml determined by ELISA (QC Test).

SPR Data



Human BAFFR, His Tag captured on CM5 Chip via anti-his antibody can bind Human BAFF (Trimer) , hFc-Flag Tag with an affinity constant of 1.21 nM as determined in SPR assay (Biacore T200).