# Cynomolgus BDCA-2 Protein

Cat. No. BCA-CM102



Description	
Source	Recombinant Cynomolgus BDCA-2 Protein is expressed from HEK293 with His tag at the N-Terminus.
	It contains Tyr48-Ile212.
Accession	A0A2K5UWP4
Molecular Weight	The protein has a predicted MW of 20.55 kDa. Due to glycosylation, the protein migrates to 28-38 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE

### Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS, 300mM L-arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please

# **Background**

BDCA-2, BDCA-3, and BDCA-4. In blood, BDCA-2 and BDCA-4 are expressed on CD11c(-) CD123(bright) plasmacytoid dendritic cells, whereas BDCA-3 is expressed on small population of CD11c() CD123(-) dendritic cells. All three Ags are not detectable on a third blood dendritic cell population, which is CD1c() CD11c(bright) CD123(dim), or on any other cells in blood.BDCA-2 is completely down-regulated on plasmacytoid CD11c(-) CD123(bright) dendritic cells, BDCA-2 is rapidly internalized at 37 degrees C after mAb labeling.

### **Assay Data**

#### Tris-Bis PAGE




Cynomolgus BDCA-2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

# **ELISA Data**

Cynomolgus BDCA-2, His Tag ELISA 0.05µg Cynomolgus BDCA-2, His Tag Per Well

minimize freeze-thaw cycles.



Immobilized Cynomolgus BDCA-2, His Tag at  $0.5\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Anti-BDCA-2 Antibody, hFc Tag with the EC50 of 5.4ng/ml determined by ELISA.