

# Human BCMA/TNFRSF17 Protein

Cat. No. BCM-HM117

## Description

<b>Source</b>	Recombinant Human BCMA/TNFRSF17 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Met1-Ala54.
<b>Accession</b>	Q02223-1
<b>Molecular Weight</b>	The protein has a predicted MW of 8.9 kDa. Due to glycosylation, the protein migrates to 12 kDa and 15-17 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1 EU per µg by the LAL method.
<b>Purity</b>	>95% as determined by Bis-Tris PAGE

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

B-cell maturation antigen (BCMA or BCM), also known as tumor necrosis factor receptor superfamily member 17 (TNFRSF17), is a protein that in humans is encoded by the TNFRSF17 gene. TNFRSF17 is a cell surface receptor of the TNF receptor superfamily which recognizes B-cell activating factor (BAFF).

## Assay Data


### Bis-Tris PAGE



Human BCMA on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### ELISA Data

**Human BCMA, His Tag ELISA**  
0.1µg Human BCMA, His Tag Per Well




Immobilized Human BCMA, His Tag at 1 µg/ml (100 µl/well) on the plate. Dose response curve for Biotinylated Human BAFF, His Tag with the EC50 of 10.1 ng/ml determined by ELISA (QC Test).

Assay Data

ELISA Data

**Human BCMA, His Tag ELISA**

0.05µg Human BCMA, His Tag Per Well



Immobilized Human BCMA, His Tag at 0.5 µg/ml (100 µl/Well) on the plate. Dose response curve for Anti-BCMA Antibody, hFc Tag with the EC50 of 2.4 ng/ml determined by ELISA.