

Human BCMA/TNFRSF17 Protein, Ultra Low Endotoxin



Cat. No. BCM-HM217-UL

Description	
Source	Recombinant Human BCMA/TNFRSF17 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Met1-Ala54.
Accession	Q02223-1
Molecular Weight	The protein has a predicted MW of 31.4 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
B-cell maturation antigen (BCMA or BCM), also known as tumor necrosis factor receptor superfamily member 17 (TNFRSF17), is a protein that in humans is encoded by the TNFRSF17 gene. TNFRSF17 is a cell surface receptor of the TNF receptor superfamily which recognizes B-cell activating factor (BAFF).	

Assay Data


Bis-Tris PAGE



Human BCMA on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human BCMA is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human BCMA, hFc Tag ELISA

0.01µg Human APRIL Trimer, His Tag Per Well



Immobilized Human APRIL Trimer, His Tag at 0.1µg/ml (100µl/well) on the plate. Dose response curve for Human BCMA, hFc Tag with the EC50 of 0.9ng/ml determined by ELISA (QC Test).