Cynomolgus BST1 Protein

Cat. No. BST-CM101

ϗνωτυς

| Description | |
|-------------------------|---|
| Source | Recombinant Cynomolgus BST1 Protein is expressed from HEK293 with His tag at the C-Terminus. |
| | It contains Gly26-Ala289. |
| Accession | A0A2K5VGB5 |
| Molecular Weight | The protein has a predicted MW of 30.99 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |
| | > 95% as determined by HPLC |
| Formulation and Storage | |
| Formulation | Supplied as 0.22µm filtered solution in PBS (pH 7.4). |
| Storage | Valid for 12 months from date of receipt when stored at -80°C.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Background | |
| | BST1 overexpression conferred resistance to sphingosine in yeast. BST1 deletion produced sensitivity to exogenous D-erythro-sphingosine and phytosphingosine and intracellular accumulation of sphingosine 1-phosphate upon exposure to exogenous sphingosine. sphingoid base metabolism is similar in all eukaryotes and suggests that yeast genetics may be useful in the isolation and identification of other genes involved in |

sphingolipid signaling and metabolism.

Assay Data





Cynomolgus BST1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cynomolgus BST1 Protein

Cat. No. BST-CM101

Assay Data





The purity of Cynomolgus BST1 is greater than 95% as determined by SEC-HPLC.