

Cynomolgus BTN3A1/CD277 Protein

Cat. No. BTN-CM4A1



Description

Source	Recombinant Cynomolgus BTN3A1/CD277 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gln1-Trp219.
Accession	A0A330KWZ1
Molecular Weight	The protein has a predicted MW of 26.5 kDa. Due to glycosylation, the protein migrates to 30-35 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The three butyrophilin BTN3A molecules, BTN3A1, BTN3A2, and BTN3A3, are members of the B7/butyrophilin-like group of Ig superfamily receptors, which modulate the function of T cells. BTN3A1 controls activation of human Vγ9/Vδ2 T cells by direct or indirect presentation of self and nonself phosphoantigens (pAg).

Assay Data


Bis-Tris PAGE



Cynomolgus BTN3A1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Cynomolgus BTN3A1 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Cynomolgus BTN3A1, His Tag ELISA

0.2 μ g Cynomolgus BTN3A1, His Tag Per Well



Immobilized Cynomolgus BTN3A1, His Tag at 2 μ g/ml (100 μ l/Well) on the plate. Dose response curve for Anti-BTN3A1 Antibody, hFc Tag with the EC50 of 6.7ng/ml determined by ELISA.