
## Cynomolgus NCAM-1/CD56 Protein





Description	
Source	Recombinant Cynomolgus NCAM-1/CD56 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Leu20-Gly718.
Accession	XP_005579710.1
Molecular Weight	The protein has a predicted MW of 78.46 kDa. Due to glycosylation, the protein migrates to 95-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


· omalaion and olorage	
Formulation	Lyophilized from 0.22 $\mu$ m filtered solution in 20mM Tris, 150mM NaCl (pH 8.0). Normally 8% mannitol is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### **Background**

Neural Cell Adhesion Molecule 1 (NCAM-1), a multifunctional member of the immunoglobulin superfamily, is expressed on the surface of neurons, glia, skeletal muscle, and natural killer cells. NCAM-1 has been implicated as having a role in cell-cell adhesion, involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance.


#### **Assay Data**

#### **Bis-Tris PAGE**



Cynomolgus NCAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### **SEC-HPLC**




The purity of Cynomolgus NCAM-1 is greater than 95% as determined by SEC-HPLC.

# KAGTUS

#### **Assay Data**

#### **ELISA Data**

# **Cynomolgus NCAM-1, His Tag ELISA** 0.05µg Cynomolgus NCAM-1, His Tag Per Well



Immobilized Cynomolgus NCAM-1, His Tag at  $0.5\mu g/ml$  (100 $\mu$ l/well) on the plate. Dose response curve for Anti-NCAM-1 Antibody, hFc Tag with the EC50 of 4.2ng/ml determined by ELISA (QC Test).