

Cynomolgus EpCAM/TROP1 Protein

Cat. No. CAM-CM1EP

Description

Source	Recombinant Cynomolgus EpCAM/TROP1 Protein is expressed from Expi293 with His tag at the C-terminal. It contains Gln24-Lys265.
Accession	NP_001035118.1
Molecular Weight	The protein has a predicted MW of 28.5 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended (usually we use 1mg/ml solution for lyophilization). Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please avoid freeze-thaw cycles.

Background

Epithelial Cellular Adhesion Molecule (EpCAM), also known as KS1/4, gp40, GA733-2, 17-1A, and TROP1, is a 40 kDa transmembrane glycoprotein that consists of a 242 amino acid (aa) extracellular domain with two EGFl like repeats, a 23 aa transmembrane segment, and a 26 aa cytoplasmic domain.

Assay Data


Tris-Bis PAGE



Cynomolgus EpCAM on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Assay Data



The purity of Cynomolgus EpCAM is greater than 95% as determined by SEC-HPLC.

SPR Data



Cynomolgus EpCAM, His Tag captured on CM5 Chip via Anti-His Antibody can bind Anti-EpCAM Antibody, hFc Tag with an affinity constant of 21.40 nM as determined in SPR assay (Biacore T200).