

Human NCAM-1/CD56 Protein

Cat. No. CAM-HM156

Description

Source	Recombinant Human NCAM-1/CD56 Protein is expressed from HEK293 with His tag at the C-Terminus It contains Leu20-Gly718.
Accession	P13591-2
Molecular Weight	The protein has a predicted MW of 78.48 kDa. Due to glycosylation, the protein migrates to 95-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Neural Cell Adhesion Molecule 1 (NCAM-1), a multifunctional member of the immunoglobulin superfamily, is expressed on the surface of neurons, glia, skeletal muscle, and natural killer cells. NCAM-1 has been implicated as having a role in cell-cell adhesion, involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance.


Assay Data

Bis-Tris PAGE



Human NCAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human NCAM-1 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human NCAM-1, His Tag ELISA

0.1µg Human NCAM-1, His Tag Per Well



Immobilized Human NCAM-1, His Tag at 1 µg/ml (100 µl/well) on the plate. Dose response curve for Anti-NCAM-1 Antibody, hFc Tag with the EC50 of 5.1 ng/ml determined by ELISA.