

Human CEACAM-5/CD66e (145-322) Protein

Cat. No. CAM-HM1D1

Description

| | |
|-------------------------|--|
| Source | Recombinant Human CEACAM-5/CD66e (145-322) Protein is expressed from HEK293 with His tag at the C-terminus It contains Pro145-Pro322. |
| Accession | P06731-1 |
| Molecular Weight | The protein has a predicted MW of 20.66 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1 EU per μg by the LAL method. |
| Purity | >95% as determined by Bis-Tris PAGE; >95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5) was identified as a metastatic driver. CEACAM5 overproduction enriched for an epithelial gene expression pattern and facilitated tumor outgrowth at metastatic sites. Tissues from patients with metastatic breast cancer confirmed elevated levels of CEACAM5 in lung metastases relative to breast tumors, and an inverse correlation between CEACAM5 and the mesenchymal marker vimentin was demonstrated.

Assay Data


Bis-Tris PAGE



Human CEACAM-5 (145-322) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human CEACAM-5 (145-322) is greater than 95% as determined by SEC-HPLC.