Human CEACAM-5/CD66e (323-500) Protein, Ultra Low Endotoxin


Cat. No. CAM-HM1D2-UL



Recombinant Human CEACAM-5/CD66e (323-500) Protein is expressed from HEK293 with His tag at the C- terminus. It contains Pro323-Leu500. P06731-1
terminus. It contains Pro323-Leu500.
P06731-1
The protein has a predicted MW of 20.73 kDa. Due to glycosylation, the protein migrates to 45-60 kDa based on Bis-Tris PAGE result.
Less than 0.01 EU per μg by the LAL method.
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC
je
Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before Iyophilization.
Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5) was identified as a metastatic driver. CEACAM5 overproduction enriched for an epithelial gene expression pattern and facilitated tumor outgrowth at metastatic sites. Tissues from patients with metastatic breast cancer confirmed elevated levels of CEACAM5 in lung metastases relative to breast tumors, and an inverse correlation between CEACAM5 and the mesenchymal marker vimentin was demonstrated.

Assay Data

Bis-Tris PAGE




Human CEACAM-5 (323-500) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Human CEACAM-5/CD66e (323-500) Protein, Ultra Low Endotoxin

Cat. No. CAM-HM1D2-UL

Assay Data





The purity of Human CEACAM-5 (323-500) is greater than 95% as determined by SEC-HPLC.