Biotinylated Human CEACAM-8/CD66b Protein





Description	
Source	Recombinant Biotinylated Human CEACAM-8/CD66b Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gln35-Ser319.
Accession	NP_001807.2
Molecular Weight	The protein has a predicted MW of 34.3 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	l Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Eosinophils and their products are likely important in the pathophysiology of allergic diseases, such as bronchial asthma, and in host immunity to parasitic organisms. CD66b (CEACAM8, CGM6, NCA-95) is a single chain, GPIanchored, highly glycosylated protein belonging to the carcinoembryonic Ag supergene family. CD66b is an activation marker for human granulocytes.

Assay Data

Tris-Bis PAGE




Biotinylated Human CEACAM-8 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. CAM-HM408B



Assay Data



The purity of Biotinylated Human CEACAM-8 is greater than 95% as determined by SEC-HPLC.


Biotinylated Human CEACAM-8/CD66b Protein

Cat. No. CAM-HM408B



Assay Data

SPR Data



Biotinylated Human CEACAM-8, His Tag immobilized on CM5 Chip can bind Human CEACAM-6, His Tag with an affinity constant of 6.13 μ M as determined in SPR assay (Biacore T200).