

Human PTK7/CCK4 Protein

Cat. No. CCK-HM104



Description

Source	Recombinant Human PTK7/CCK4 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Ala31-Thr704.
Accession	Q13308-1
Molecular Weight	The protein has a predicted MW of 75.7 kDa. Due to glycosylation, the protein migrates to 80-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Protein Tyrosine Kinase 7 (PTK7) is a critical regulator of canonical and non-canonical Wnt-signaling during embryonic development and cancer cell formation. Disrupting PTK7 activity perturbs vertebrate nervous system development, and also promotes human cancer formation. Observations in different model systems suggest a complex cross-talk between PTK7 protein and Wnt signaling.


Assay Data

Bis-Tris PAGE



Human PTK7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human PTK7 is greater than 95% as determined by SEC-HPLC.