
#### Biotinylated Human CD155/PVR Protein






Description	
Source	Recombinant Biotinylated Human CD155/PVR Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Trp21-Asn343.
Accession	P15151-1
Molecular Weight	The protein has a predicted MW of 38 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Bis- Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	

# CD155 is a cell surface adhesion molecule functioning in tumor cell migration, invasion, and metastasis, and not surprisingly, is also designated as a common tumor-associated antigen. CD155 is also recognized by NK cells to induce their cytotoxicity. CD155 is also commonly referred to as the "poliovirus receptor," or PVR.


#### **Assay Data**

#### **Bis-Tris PAGE**



Biotinylated Human CD155 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

#### SEC-HPLC



The purity of Biotinylated Human CD155 is greater than 95% as determined by SEC-HPLC.

## KAGTUS

#### **Assay Data**

#### **ELISA Data**


## **Biotinylated Human CD155, His Tag ELISA** 0.1µg Biotinylated Human CD155, His Tag Per Well

3.0 (05400) square 1.0 0.5 0.0 -6 -5 -4 -3 -2 -1 0 Log Human TIGIT, hFc Tag Conc.(µg/ml)

Immobilized Biotinylated Human CD155, His Tag at  $1\mu g/ml$  (100 $\mu l/well$ ) on the streptavidin precoated plate ( $5\mu g/ml$ ). Dose response curve for Human TIGIT, hFc Tag with the EC50 of 3.0ng/ml determined by ELISA (QC Test).

#### **ELISA Data**

## **Biotinylated Human CD155, His Tag ELISA** 0.1μg Biotinylated Human CD155, His Tag Per Well



Immobilized Biotinylated Human CD155, His Tag at  $1\mu g/ml$  ( $100\mu l/well$ ) on the plate. Dose response curve for Human DNAM-1, hFc Tag with the EC50 of 10.6ng/ml determined by ELISA.