Mouse IL-2 R beta/CD122 Protein

Cat. No. CD1-MM122

Description

Source	Recombinant Mouse IL-2 R beta/CD122 Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Ala27-Glu240.
Accession	P16297
Molecular Weight	The protein has a predicted MW of 26.13 kDa. Due to glycosylation, the protein migrates to 50-65 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IL-2 R beta is a member of the cytokine receptor superfamily. Human IL-2 R beta cDNA encodes a 551 amino acid residue precursor Type I membrane protein with a 26 residue signal peptide, a 214 residue extracellular region, a 25 residue transmembrane region and a 286 residue cytoplasmic domain. Functional IL-2 receptors can exist in two affinity states on cell surfaces, the high affinity complex consisting of heterotrimers of the alpha, beta, and gamma chains, and the intermediate affinity complex comprising heterodimers of the beta and gamma chains.

Assay Data

Bis-Tris PAGE




Mouse IL-2 R beta on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Mouse IL-2 R beta/CD122 Protein

Cat. No. CD1-MM122

Assay Data




The purity of Mouse IL-2 R beta is greater than 95% as determined by SEC-HPLC.

Mouse IL-2 R beta/CD122 Protein

Cat. No. CD1-MM122

Assay Data

SPR Data



Mouse IL-2 R beta, His Tag captured on CM5 Chip via Anti-His Antibody can bind Human IL-2, No Tag (Cat. IL2-HM001) with an affinity constant of 2.39 μM as determined in SPR assay (Biacore T200).

For Research Use Only