
## Cynomolgus/Rhesus macaque CD27/TNFRSF7 Protein





Description	
Source	Recombinant Cynomolgus/Rhesus macaque CD27/TNFRSF7 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Thr21-Ile192.
Accession	XP_005569963.1(Cynomolgus) / F7BYS2(Rhesus macaque)
Molecular Weight	The protein has a predicted MW of 20.41 kDa. Due to glycosylation, the protein migrates to 42-52 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### Background

CD27, also known as TNFRSF7, is an approximately 55 kDa transmembrane protein in the TNF receptor superfamily. It functions as a costimulatory molecule that supports lymphocyte activation and survival. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis.


#### **Assay Data**

#### **Bis-Tris PAGE**



Cynomolgus/Rhesus macaque CD27 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC



The purity of Cynomolgus/Rhesus macaque CD27 is greater than 95% as determined by SEC-HPLC.


Cat. No. CD2-CM127



#### **Assay Data**

**ELISA Data** 

# **Cynomolgus CD27, His Tag ELISA** 0.1μg Cynomolgus CD27, His Tag Per Well



Immobilized Cynomolgus/Rhesus macaque CD27, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human CD27 Ligand, hFc Tag with the EC50 of 17.6ng/ml determined by ELISA.