

Mouse CD27/TNFRSF7 Protein

Cat. No. CD2-MM327



Description

Source	Recombinant Mouse CD27/TNFRSF7 Protein is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus. It contains Pro24-Arg182.
Accession	P41272-1
Molecular Weight	The protein has a predicted MW of 44.3 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD27, also known as TNFRSF7, is an approximately 55 kDa transmembrane protein in the TNF receptor superfamily. It functions as a costimulatory molecule that supports lymphocyte activation and survival. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis.


Assay Data

Bis-Tris PAGE



Mouse CD27 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Mouse CD27 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Mouse CD27, mFc Tag ELISA

0.2µg Mouse CD27, mFc Tag Per Well



Immobilized Mouse CD27, mFc Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Mouse CD27 Ligand, hFc Tag with the EC50 of 34.0ng/ml determined by ELISA (QC Test).