

Biotinylated Cynomolgus CD3E/CD3 epsilon 1-27 Protein

Cat. No. CD3-CM2EDB

Description

Source	Recombinant Biotinylated Cynomolgus CD3E/CD3 epsilon 1-27 Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus. It contains Asp22-Thr48.
Accession	Q95LI5.2
Molecular Weight	The protein has a predicted MW of 31.3 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD3E, is a single-pass type I membrane protein. CD3 (cluster of differentiation 3) T cell co-receptor helps to activate both the cytotoxic T cell (CD8 naive T cells) and also T helper cells (CD4 naive T cells). It consists of a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3γ chain, a CD3δ chain, and two CD3ε chains.


Assay Data

Bis-Tris PAGE



Biotinylated Cynomolgus CD3E 1-27 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Cynomolgus CD3E 1-27 is greater than 95% as determined by SEC-HPLC.

Biotinylated Cynomolgus CD3E/CD3 epsilon 1-27 Protein

Cat. No. CD3-CM2EDB




Assay Data

ELISA Data

Biotinylated Cynomolgus CD3E 1-27, hFc Tag ELISA

0.1µg Anti-CD3 Antibody, hFc Tag Per Well



Immobilized Anti-CD3 Antibody, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus CD3E 1-27, hFc Tag with the EC50 of 5.8ng/ml determined by ELISA.

Log Biotinylated Cynomolgus CD3E 1-27, hFc Tag Conc.(µg/ml)