Biotinylated Human CD23/Fc epsilon RII Protein


Cat. No. CD3-HM123B



Description	
Source	Recombinant Biotinylated Human CD23/Fc epsilon RII Protein is expressed from HEK293 with His tag and Avitag at the N-terminus.
	It contains Asp48-Ser321.
Accession	P06734-1
Molecular Weight	The protein has a predicted MW of 33.89 kDa. Due to glycosylation, the protein migrates to 42-52 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and St	torage
Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	CD23 is the low-affinity receptor for immunoglobulin (Ig)E and plays important roles in the regulation of IgE responses. CD23 can be cleaved from cell surfaces to yield a range of soluble CD23 (sCD23) proteins that have pleiotropic cytokine-like activities.

Assay Data

Bis-Tris PAGE




10KD SEC-HPLC Biotinylated Human CD23 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Cat. No. CD3-HM123B



Assay Data



The purity of Biotinylated Human CD23 is greater than 95% as determined by SEC-HPLC.