
Biotinylated Human CD38 Protein





Description	
Source	Recombinant Biotinylated Human CD38 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Val43-Ile300.
Accession	P28907-1
Molecular Weight	The protein has a predicted MW of 32.50 kDa. Due to glycosylation, the protein migrates to 42-50 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please

Background

CD38 (cluster of differentiation 38), also known as cyclic ADP ribose hydrolase, is a transmembrane glycoprotein found on the surface of some immune cells including plasma cells, activated or immature T and B cells, monocytes, and natural killer cells. CD38 participates in cell adhesion, signal transduction and calcium signaling.


Assay Data

Tris-Bis PAGE



Biotinylated Human CD38 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human CD38 is greater than 95% as determined by SEC-HPLC.




Assay Data

ELISA Data

Biotinylated Human CD38, His Tag ELISA

0.1μg Biotinylated Human CD38, His Tag Per Well



Log Anti-CD38 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Biotinylated Human CD38, His Tag at $1\mu g/ml$ ($100\mu l/well$) on the streptavidin precoated plate($5\mu g/ml$). Dose response curve for Anti-CD38 Antibody, hFc Tag with the EC50 of 8.1ng/ml determined by ELISA.