Human CD30/TNFRSF8 Protein

Cat. No. CD3-HM430



Description	
Source	Recombinant Human CD30/TNFRSF8 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Phe19-Lys379.
Accession	P28908-1
Molecular Weight	The protein has a predicted MW of 41.3 kDa. Due to glycosylation, the protein migrates to 68-95 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	1 Storago

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The transmembrane receptor CD30 (TNFRSF8) and its ligand CD30L (CD153, TNFSF8) are members of the tumor necrosis factor (TNF) superfamily and display restricted expression in subpopulations of activated T-and B-cells in nonpathologic conditions. CD30 expression is upregulated in various hematological malignancies, including Reed-Sternberg cells in Hodgkin's disease (HD), anaplastic large cell lymphoma (ALCL) and subsets of Non-Hodgkin's lymphomas (NHLs).


Assay Data

Bis-Tris PAGE



Human CD30 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human CD30 is greater than 95% as determined by SEC-HPLC.

Human CD30/TNFRSF8 Protein


Cat. No. CD3-HM430

KAGTUS

Assay Data

ELISA Data

Human CD30, His Tag ELISA 0.05μg Human CD30, His Tag Per Well



Immobilized Human CD30, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Human CD30 Ligand, mFc Tag with the EC50 of 21.4ng/ml determined by ELISA.