

Biotinylated Human CD3E&CD3D/CD3 epsilon&CD3 delta Protein

Cat. No. CD3-HM505B

Description

Source	Recombinant Biotinylated Human CD3E&CD3D/CD3 epsilon&CD3 delta Protein is expressed from Expi293 with hFc tag and Avi tag at the C-terminal. It contains Asp23-Asp126(CD3E) & Phe22-Ala105(CD3D).
Accession	P07766(CD3E)&P04234(CD3D)
Molecular Weight	The protein has a predicted MW of 40 kDa (CD3E) and 35.4 kDa (CD3D). Due to glycosylation, the protein migrates to 48-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please avoid freeze-thaw cycles.

Background

T-cell surface glycoprotein CD3 epsilon&CD3 delta chain, also known as CD3E&CD3D, are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.

Assay Data


Tris-Bis PAGE



Biotinylated Human CD3E&CD3D on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Biotinylated Human CD3E&CD3D was greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human CD3E&CD3D, hFc Tag ELISA

0.1 µg Biotinylated Human CD3E&CD3D, hFc Tag Per Well



Immobilized Biotinylated Human CD3E&CD3D, hFc Tag at 1 µg/ml (100 µl/Well) on streptavidin (5 µg/ml) precoated plate. Dose response curve for OKT3, mFc Tag with the EC50 of 27.7 ng/ml determined by ELISA.