

Cynomolgus IL-2 R alpha/CD25 Protein

Cat. No. CD5-CM125



Description

Source	Recombinant Cynomolgus IL-2 R alpha/CD25 Protein is expressed from HEK293 with His tag at the C-terminus. It contains Glu22-Glu238.
Accession	H6WS54
Molecular Weight	The protein has a predicted MW of 69.61 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Supplied as 0.22 μm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C . Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IL-2 receptor alpha (IL-2 R alpha), also known as Tac antigen and as CD25, was initially identified as a 55 kDa membrane glycoprotein that is capable of binding IL2.eceptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.


Assay Data

Bis-Tris PAGE



Cynomolgus IL-2 R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Cynomolgus IL-2 R alpha is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Cynomolgus IL-2 R alpha, His Tag ELISA

0.5µg Cynomolgus IL-2 R alpha, His Tag Per Well



Immobilized Cynomolgus IL-2 R alpha, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human IL-2, His Avi Tag (Cat. IL2-HM401B) with the EC50 of 0.15µg/ml determined by ELISA.