
## Biotinylated Human IL-2 R alpha/CD25 Protein





Description	
Source	Recombinant Biotinylated Human IL-2 R alpha/CD25 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Glu22-Cys213.
Accession	P01589
Molecular Weight	The protein has a predicted MW of 24.6 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

## Formulation and Storage


Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

Human IL-2 receptor alpha (IL-2 R alpha), also known as Tac antigen and as CD25, was initially identified as a 55 kDa membrane glycoprotein that is capable of binding IL2.eceptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.


### **Assay Data**

### **Bis-Tris PAGE**



Biotinylated Human IL-2 R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### **SEC-HPLC**



The purity of Biotinylated Human IL-2 R alpha is greater than 95% as determined by SEC-HPLC.




### **Assay Data**

### **ELISA Data**

# Biotinylated Human IL-2 R alpha, His Avi Tag ELISA

0.05μg Biotinylated Human IL-2 R alpha, His Avi Tag Per Well



Log Anti-IL-2 R alpha Antibody, hFc Tag Conc.(μg/ml)

Immobilized Biotinylated Human IL-2 R alpha, His Avi Tag at  $0.5\mu g/ml$  (100 $\mu l/well$ ) on the streptavidin precoated plate (5 $\mu g/ml$ ). Dose response curve for Anti-IL-2 R alpha Antibody, hFc Tag with the EC50 of 3.9 ng/ml determined by ELISA (QC Test).