
Human CD69/CLEC2C Protein






Cat. 140. OD0-11101150	•
Description	
Source	Recombinant Human CD69/CLEC2C Protein is expressed from HEK293 with His tag at the N-Terminus.
	It contains Ser62-Lys199.
Accession	Q07108
Molecular Weight	The protein has a predicted MW of 17.07kDa. Due to glycosylation, the protein migrates to 24-27kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	CLEC2C (CD69) is a membrane-bound, type II C-lectin receptor and acts as a costimulatory molecule for T cell activation and proliferation. It is involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. CLEC2C is a disulfide-linked homodimer protein with two differentially glycosylated subunits.

Assay Data

Bis-Tris PAGE



Human CD69 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.