

Human IL-17R alpha/CD217 Protein

Cat. No. CD7-HM2RA

Description

Source	Recombinant Human IL-17R alpha/CD217 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Leu33-Trp320.
Accession	Q96F46-1
Molecular Weight	The protein has a predicted MW of 60.2 kDa. Due to glycosylation, the protein migrates to 75-90 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin 17 (also known as CTLA8) is a T cell-expressed pleiotropic cytokine. IL17 binds to IL17 receptor (IL17 R) which shares no homology with any known family of receptors. While the expression of IL17 is restricted to activated T cells, the IL17 R mRNA exhibits a broad tissue distribution, and has been detected in virtually all cells and tissues tested. The human IL17 R gene was localized to chromosome 22.


Assay Data

Bis-Tris PAGE



Human IL-17R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human IL-17R alpha is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human IL-17R alpha, hFc Tag ELISA

0.1µg Human IL-17A, His Tag Per Well




Immobilized Human IL-17A, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human IL-17R alpha, hFc Tag with the EC50 of 72.2ng/ml determined by ELISA.

ELISA Data

Human IL-17R alpha, hFc Tag ELISA

0.5µg Human IL-17F, His Tag Per Well




Immobilized Human IL-17F, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Human IL-17 R alpha, hFc Tag with the EC50 of 1.75µg/ml determined by ELISA.

ELISA Data

Human IL-17R alpha, hFc Tag ELISA


0.1µg Human IL-17A&F, His Tag Per Well



Immobilized Human IL-17A&F, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human IL-17R alpha, hFc Tag with the EC50 of 44.1ng/ml determined by ELISA.

SPR Data

Assay Data



Human IL-17R alpha, hFc Tag captured on CM5 Chip via protein A can bind Biotinylated Human IL-17A, His-Avi Tag with an affinity constant of 2.18 nM as determined in SPR assay (Biacore T200).