

Human CD93/C1q R1 Protein

Cat. No. CD9-HM193

Description

Source	Recombinant Human CD93/C1q R1 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Thr22-Lys580.
Accession	Q9NPY3
Molecular Weight	The protein has a predicted MW of 59.4 kDa. Due to glycosylation, the protein migrates to 95-110 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD93 has been shown critical roles in inflammatory and immune diseases. CD93 silencing increased IL-6 and TSLP, but not IL-33 levels in culture supernatants. HDM-induced asthma mice showed significant airway hyperresponsiveness and inflammation with Th2 cytokine activation, along with decreased CD93 expression in bronchial epithelial cells and lung homogenates but increased serum CD93 levels.

Assay Data


Bis-Tris PAGE



Human CD93 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human CD93 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human CD93, His Tag ELISA

0.5µg Human CD93, His Tag Per Well



Immobilized Human CD93, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Human IGFBP-7, hFc Tag with the EC50 of 0.69µg/ml determined by ELISA (QC Test).