

Human CD99/MIC2 Protein

Cat. No. CD9-HM299

Description

Source	Recombinant Human CD99/MIC2 Protein is expressed from Expi293 with hFc tag at the C-terminal. It contains Asp23-Asp122.
Accession	AAH02584
Molecular Weight	The protein has a predicted MW of 36.7 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended (usually we use 1mg/ml solution for lyophilization). Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please avoid freeze-thaw cycles.

Background

CD99 is a cell surface protein with unique features and only partly defined mechanisms of action. This molecule is involved in crucial biological processes, including cell adhesion, migration, death, differentiation and diapedesis, and it influences processes associated with inflammation, immune responses and cancer. CD99 is frequently overexpressed in many types of tumors, particularly pediatric tumors including Ewing sarcoma and specific subtypes of leukemia.

Assay Data


Tris-Bis PAGE



Human CD99 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human CD99 is greater than 95% as determined by SEC-HPLC.