### Human CD96/TACTILE Protein

#### Cat. No. CD9-HM396


# ᠺ᠕ᢏᠴᡃ᠐᠍᠋ᠶ

Description Recombinant Human CD96/TACTILE Protein is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus. Source It contains Val22-Met503. Accession P40200-2 The protein has a predicted MW of 79.8 kDa. Due to glycosylation, the protein migrates to 120-160 kDa based on Molecular Weight Bis-Tris PAGE result. Endotoxin Less than 1EU per µg by the LAL method. > 95% as determined by Bis-Tris PAGE Purity > 95% as determined by HPLC Formulation and Storage Formulation Supplied as 0.22µm filtered solution in PBS (pH 7.4). Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller Storage quantities for optimal storage. Please minimize freeze-thaw cycles. Background The receptors CD96 and TIGIT are expressed on the surface of T and natural killer (NK) cells, and recent studies suggest both play important inhibitory roles in immune function. CD96 has been shown to modulate immune cell activity in mice, with Cd96-/- mice displaying hypersensitive NK-cell responses to immune challenge and significant tumor resistance. The counterbalance between the putative inhibitory CD96 and TIGIT receptors and

the activating receptor, CD226, offers unique strategies for immuno-oncology drug development.

Assay Data

**Bis-Tris PAGE** 




Human CD96 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Human CD96, mFc Tag ELISA

0.5μg Human CD96, mFc Tag Per Well



Immobilized Human CD96, mFc Tag at  $5\mu$ g/ml (100 $\mu$ I/Well) on the plate. Dose response curve for Human CD155, hFc Tag with the EC50 of 0.19 $\mu$ g/ml determined by ELISA (QC Test).


# Human CD96/TACTILE Protein

Cat. No. CD9-HM396

### **Assay Data**

## ELISA Data

Human CD96, mFc Tag ELISA 0.2µg Human CD96, mFc Tag Per Well



Immobilized Human CD96, mFc Tag at  $2\mu$ g/ml (100 $\mu$ I/Well) on the plate. Dose response curve for Anti-CD96 Antibody, hFc Tag with the EC50 of 31.8ng/ml determined by ELISA.

