

Human CD300A Protein

Cat. No. CDA-HM101



Description

| | |
|-------------------------|--|
| Source | Recombinant Human CD300A Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Leu18-Pro180. |
| Accession | Q9UGN4-1 |
| Molecular Weight | The protein has a predicted MW of 18.7 kDa. Due to glycosylation, the protein migrates to 48-60 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

CD300a is an inhibitory receptor for mast cells and eosinophils in allergic inflammation (AI); however, the spatiotemporal expression of CD300a and its potential roles in the resolution of AI are still to be determined. CD300a expression on peritoneal cells is regulated from inflammation to resolution. CD300a activation on mouse bone marrow-derived mast cells regulated ALX/FPR2 expression levels following IgE-mediated activation.


Assay Data

Bis-Tris PAGE



Human CD300A on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human CD300A is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human CD300A, His Tag ELISA

0.2µg Human CD300A, His Tag Per Well



Immobilized Human CD300A, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-CD300A Antibody, hFc Tag with the EC50 of 3.6ng/ml determined by ELISA.