

Cynomolgus Fc gamma RIIB Protein

Cat. No. CDB-CM101



Description

Source	Recombinant Cynomolgus Fc gamma RIIB Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ala46-Pro224.
Accession	Q8SPW3
Molecular Weight	The protein has a predicted MW of 21 kDa. Due to glycosylation, the protein migrates to 33-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Supplied as 0.22 μm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. The CD64 proteins are high affinity receptors ($\sim 10\text{e-}8$ - $10\text{e-}9$ M) capable of binding monomeric IgG, whereas the CD16 and CD32 proteins bind IgG with lower affinities ($\sim 10\text{e-}6$ - $10\text{e-}7$ M) only recognizing IgG aggregates surrounding multivalent antigens.


Assay Data

Bis-Tris PAGE



Cynomolgus Fc gamma RIIB on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC



The purity of Cynomolgus Fc gamma RIIB is greater than 95% as determined by SEC-HPLC.

Assay Data

SPR Data



Cynomolgus Fc gamma RIIB, His Tag captured on CM5 Chip via Anti-His Antibody can bind Trastuzumab with an affinity constant of 1.75 μ M as determined in SPR assay (Biacore T200).