Human CDCP1 (30-368) Protein

Cat. No. CDC-HM10D



Description	
Source	Recombinant Human CDCP1 (30-368) Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Phe30-Arg368.
Accession	Q9H5V8-1
Molecular Weight	The protein has a predicted MW of 39.03 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Tumor metastasis depends on the dynamic regulation of cell adhesion through β1-integrin. The Cub-Domain Containing Protein-1, CDCP1, is a transmembrane glycoprotein which regulates cell adhesion. Overexpression and loss of CDCP1 have been observed in the same cancer types to promote metastatic progression.


Assay Data

Bis-Tris PAGE



Human CDCP1 (30-368) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human CDCP1 (30-368) is greater than 95% as determined by SEC-HPLC.

KAGTUS

Assay Data

ELISA Data

Human CDCP1 (30-368), His Tag ELISA 0.2μg Human CDCP1 (30-368), His Tag Per Well



Immobilized Human CDCP1 (30-368), His Tag at $2\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Anti-CDCP1 Antibody, hFc Tag with the EC50 of 2.8ng/ml determined by ELISA.