

Biotinylated Cynomolgus CDH17/Cadherin 17 Protein

Cat. No. CDH-CM427B

Description

Source	Recombinant Biotinylated Cynomolgus CDH17/Cadherin 17 Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus. It contains Gln23-Thr784.
Accession	XP_005563762.2
Molecular Weight	The protein has a predicted MW of 87.39 kDa. Due to glycosylation, the protein migrates to 110-130 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 90% as determined by Bis-Tris PAGE > 85% as determined by HPLC

Formulation and Storage

Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Liver-intestine cadherin (CDH17) has been known to function as a tumor stimulator and diagnostic marker for almost two decades. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.


Assay Data

Bis-Tris PAGE



Biotinylated Cynomolgus CDH17 on Bis-Tris PAGE under reduced condition. The purity is greater than 90%.

SEC-HPLC




The purity of Biotinylated Cynomolgus CDH17 is greater than 85% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Cynomolgus CDH17, His Avi Tag ELISA

0.05µg Biotinylated Cynomolgus CDH17, His Avi Tag Per Well



Immobilized Biotinylated Cynomolgus CDH17, His Avi Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-CDH17 Antibody, hFc Tag with the EC50 of 2.3ng/ml determined by ELISA.