Human E-Cadherin/Cadherin-1 Protein

Cat. No. CDH-HM101



Description	
Source	Recombinant Human E-Cadherin/Cadherin-1 Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Asp155-Ala709.
Accession	P12830-1
Molecular Weight	The protein has a predicted MW of 61.62 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 90% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 μ m filtered solution in 20mM Tris, 150mM NaCL (pH 8.0). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.


E-cadherin is the core component of epithelial adherens junctions, essential for tissue development,

differentiation, and maintenance. It is also fundamental for tissue barrier formation, a critical function of epithelial

Assay Data

Background

Bis-Tris PAGE



tissues.


Human E-Cadherin on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. CDH-HM101

KAGTUS

Assay Data



The purity of Human E-Cadherin is greater than 90% as determined by SEC-HPLC.