# Human CDH17/Cadherin 17 Protein, Ultra Low Endotoxin





| Description         |   |
|---------------------|---|
| Source              | Recombinant Human CDH17/Cadherin 17 Protein is expressed from HEK293 with His tag at the C-terminus.                                |
|                     | It contains Gln23-Met787.   |
| Accession           | Q12864-1  |
| Molecular<br>Weight | The protein has a predicted MW of 86.1 kDa. Due to glycosylation, the protein migrates to 90-115 kDa based on Bis-Tris PAGE result. |
| Endotoxin           | Less than 0.01 EU per μg by the LAL method.   |
| Purity              | > 95% as determined by Bis-Tris PAGE  |
|                     | > 95% as determined by HPLC   |
|                     |   |

## Formulation and Storage


| Formulation    | Lyophilized from 0.22 $\mu$ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.  |
|----------------|---|
| Reconstitution | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.   |
| Storage        | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

# Background

Liver-intestine cadherin (CDH17) has been known to function as a tumor stimulator and diagnostic marker for almost two decades. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.

## **Assay Data**

### **Bis-Tris PAGE**




Human CDH17 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS


# **Assay Data**



The purity of Human CDH17 is greater than 95% as determined by SEC-HPLC.


#### **ELISA Data**

# Human CDH17, His Tag ELISA 0.1µg Human CDH17, His Tag Per Well



Immobilized Human CDH17, His Tag at 1  $\mu$ g/ml (100  $\mu$ l/Well) on the plate. Dose response curve for Anti-CDH17 Antibody, hFc Tag with the EC50 of 4.1  $\mu$ g/ml determined by ELISA.

# **BLI Data**



Loaded Anti-CDH17 Ab., hFc Tag on ProA-Biosensor can bind Human CDH17, His Tag with an affinity constant of 2.26 nM as determined in BLI assay (Gator® Prime).