

Human CD44v9 Protein

Cat. No. CDX-HM2V9



Description

Source	Recombinant Human CD44v9 Protein is expressed from HEK293 with hFc (IgG1) tag at the C-terminus.
	It contains Gln21-Pro220 & Gly506-Ser534.
Accession	P16070-1
Molecular Weight	The protein has a predicted MW of 51.17 kDa. Due to glycosylation, the protein migrates to 80-100 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD44 is a hyaluronan binding cell surface signal transducing receptor that influences motility, cell survival and proliferation as well as the formation of tumor microenvironment. CD44 contains two variable regions encoded by variable exons. Alternative splicing, which is often deregulated in cancer, can produce various isoforms of CD44 with properties that may have different tissue specific effects and therefore even diverse effects on cancer progression


Assay Data

Bis-Tris PAGE



Human CD44v9 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human CD44v9 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human CD44v9, hFc Tag ELISA
0.05µg Human CD44v9, hFc Tag Per Well



Immobilized Human CD44v9, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-CD44v9 Antibody, hFc Avi Tag with the EC50 of 13.1ng/ml determined by ELISA.